

P-6149



PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re U.S. Patent Application)

Applicant: Botha et al.)

Serial No.: 09/779,237)

Filed: February 8, 2001)

For: The Regulation and)
Manipulation of Sucrose)
Content in Sugarcane)
)
)
)

CERTIFIED COPY OF PRIORITY APPLICATION

Sertifikaat



REPUBLIEK VAN SUID-AFRIKA



REPUBLIC OF SOUTH AFRICA

Certificate

PATENTKANTOOR

PATENT OFFICE

DEPARTEMENT VAN HANDEL
EN NYWERHEIDDEPARTMENT OF TRADE
AND INDUSTRYHiermee word gesertifiseer dat
This is to certify that

the documents attached hereto are true copies of the Forms P2, P6,
provisional specification and drawings of South African Patent Application No. 2000/0574 in
the name of South African Sugar Association

Filed

:

8 February 2000

Entitled

:

The Regulation and Manipulation of
Sucrose Content in Sugarcane

Geteken te
Signed at

PRETORIA

in die Republiek van Suid-Afrika, hierdie
in the Republic of South Africa, this

dag van
day of

14 Februarie 2000

Registrateur van Patente
Registrar of Patents

REPUBLIC OF SOUTH AFRICA			REGISTER OF PATENTS			PATENTS ACT, 1		
OFFICIAL APPLICATION			LODGING DATE: PROVISIONAL			ACCEPTANCE DATE		
21	01	20000574	22	8 FEBRUARY 2000		47		
INTERNATIONAL CLASSIFICATION			LODGING DATE: COMPLETE			GRANTED DATE		
51			23					
FULL NAME(S) OF APPLICANT(S)/PATENTEE(S)								
71	SOUTH AFRICAN SUGAR ASSOCIATION							
APPLICANTS SUBSTITUTED:						DATE REGISTERED		
71								
ASSIGNEE(S)						DATE REGISTERED		
71								
FULL NAME(S) OF INVENTOR(S)								
72	1. BOTHA; FREDERIK COENRAAD 2. GROENEWALD; JAN HENDRIK							
PRIORITY CLAIMED		COUNTRY		NUMBER		DATE		
N.B. Use International abbreviation for country (see Schedule 4)		33	NONE	31	NONE	32	NONE	
TITLE OF INVENTION								
54	THE REGULATION AND MANIPULATION OF SUCROSE CONTENT IN SUGARCANE							
ADDRESS OF APPLICANT(S), PATENTEE(S)								
MOUNT EDGECOMBE, KWAZULU-NATAL, SOUTH AFRICA								
ADDRESS FOR SERVICE						S AND F REF		
74	SPOOR AND FISHER, SANDTON					JP/T 551 /al		
PATENT OF ADDITION NO.				DATE OF ANY CHANGE				
61								
FRESH APPLICATION BASED ON				DATE OF ANY CHANGE				

2000/50574

0.0000

R 060,00

The granting of a patent is hereby requested by the undermentioned applicant on the basis of the present application filed in duplicate

REPUBLIC OF SOUTH AFRICA

OFFICIAL APPLICATION NO.

S AND F REFERENCE

21	01	20000574
----	----	----------

JP/T 551 /al

FULL NAME(S) OF APPLICANT(S)

71	SOUTH AFRICAN SUGAR ASSOCIATION
----	---------------------------------

ADDRESS(ES) OF APPLICANT(S)

	MOUNT EDGECOMBE, KWAZULU-NATAL, SOUTH AFRICA
--	--

TITLE OF INVENTION

54	THE REGULATION AND MANIPULATION OF SUCROSE CONTENT IN SUGARCANE
----	---

THE APPLICANT CLAIMS PRIORITY AS SET OUT ON THE ACCOMPANYING FORM P.2 THE EARLIEST PRIORITY CLAIM IS:

COUNTRY: NONE

NUMBER: NONE

DATE: NONE

THIS APPLICATION IS FOR A PATENT OF ADDITION TO PATENT APPLICATION NO.

21	01	
----	----	--

THIS APPLICATION IS A FRESH APPLICATION IN TERMS OF SECTION 37 AND IS BASED ON APPLICATION NO.


21	01	
----	----	--

THIS APPLICATION IS ACCOMPANIED BY:

- ☒ 1. A single copy of a provisional or two copies of a complete specification of 8 pages.
- ☒ 2. Drawings of 4 sheets.
- ☐ 3. Publication particulars and abstract (Form P.8 in duplicate).
- ☐ 4. A copy of Figure of the drawings (if any) for the abstract.
- ☐ 5. An assignment of invention.
- ☐ 6. Certified priority document(s).
- ☐ 7. Translation of the priority document(s).
- ☐ 8. An assignment of priority rights.
- ☐ 9. A copy of the Form P.2 and the specification of S.A. Patent Application No.
- ☐ 10. A declaration and power of attorney on Form P.3.
- ☐ 11. Request for ante-dating on Form P.4.
- ☐ 12. Request for classification on Form P.9.
- ☒ 13. Form P.2 in duplicate.

74 ADDRESS FOR SERVICE: SPOOR AND FISHER, SANDTON

Dated: 8 FEBRUARY 2000



SPOOR AND FISHER
PATENT ATTORNEYS FOR THE APPLICANT(S)

RECEIVED

2000-02-08

REGISTERED PATENT OFFICE
PATENT ATTORNEYS FOR THE APPLICANT(S)

REGISTRAR OF PATENTS

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978

PROVISIONAL SPECIFICATION

(Section 30(1) - Regulation 27)

OFFICIAL APPLICATION NO.

LODGING DATE

21	01	20000574
----	----	----------

22	8 FEBRUARY 2000
----	-----------------

FULL NAME(S) OF APPLICANT(S)

71	SOUTH AFRICAN SUGAR ASSOCIATION
----	---------------------------------

FULL NAME(S) OF INVENTOR(S)

72	1. BOTHA; FREDERIK COENRAAD • 2. GROENEWALD; JAN HENDRIK
----	---

TITLE OF INVENTION

54	THE REGULATION AND MANIPULATION OF SUCROSE CONTENT IN SUGARCANE
----	---

BACKGROUND OF THE INVENTION

THIS invention relates to the regulation and manipulation of sucrose content in sugarcane.

The amount of sucrose which accumulates in the culm of sugarcane is a function of sink strength and carbon partitioning between competing metabolic pathways.

Pyrophosphate-dependent phosphofructokinase (PFP) with the international identification of (EC 2.7.1.90) catalyses the reversible conversion of fructose 6-phosphate (F6P) and pyrophosphate (PPi) to fructose 1,6-bisphosphate and inorganic phosphate. As such, the enzyme probably plays an important role in regulating carbon partitioning, and therefore sucrose

levels, in sugarcane. The levels of both PPi and F6P control the rate of sucrose accumulation by affecting the rate of sucrose synthesis and hydrolysis. Consistent with this hypothesis, the activity of PFP in sugarcane increases with culm maturity, and varies between varieties differing in their sucrose storage capacity.

PFP is associated with sink tissues, and plays a major role in sink strength. PFP is a heterotetramer with two α and two β -subunits. The β -subunit is the catalytic subunit of the enzyme. Removal of the β -subunit will therefore reduce the PFP activity.

It is an object of the invention to use the sugarcane PFP- β gene or part thereof in an antisense and untranslatable form to regulate the levels of PFP in sugarcane.

SUMMARY OF THE INVENTION

According to the invention an isolated nucleotide sequence comprises:

- (i) a nucleotide sequence as set out in Figure 1;
- (ii) a nucleotide sequence which is complementary to the nucleotide sequence of (i);
- (iii) a variant of the nucleotide sequence of (i);
- (iv) a portion of the nucleotide sequence of (i); and
- (v) a nucleotide sequence which hybridizes to the nucleotide sequence of (i) under stringent hybridization conditions.

The nucleotide sequence may be the nucleotide sequence as set out in Figure

3.

The nucleotide sequence may be in an antisense orientation.

According to another aspect of the invention a gene construct comprises a promoter and nucleotide sequence as defined herein in a sense orientation, the gene construct lacking a translation initiation codon upstream of the nucleotide sequence or possessing an in-frame termination codon directly downstream of the initiation codon.

The gene construct may comprise two promoters.

The promoters may be selected from the CaMV35S and the maize polyubiquitin (UBI) promoters.

According to another aspect of the invention a gene construct comprises a promoter and a nucleotide sequence as defined herein in an antisense orientation.

The gene construct may comprise two promoters.

The promoters may be selected from the CaMV35S and the maize polyubiquitin (UBI) promoters.

The gene constructs may be expression vectors, pUSPC 510 and pASPC 510 respectively.

According to another aspect of the invention a transformed sugarcane plant

cell comprises a gene construct of the invention.

According to another aspect of the invention a transgenic sugarcane plant or sugarcane plant part containing or derived from the plant cell is provided.

The transgenic sugarcane plant part may be a sugarcane callus.

The transformed sugarcane cell or transgenic plant or plant part may be characterized by a lower level of the PFP β -subunit.

The transformed sugarcane cell or transgenic plant or plant part may be characterized by a higher level of sucrose.

According to another aspect of the invention a method of regulating or manipulating the level of active PFP in a plant cell comprises the step of transforming the plant cell with at least one gene construct of the invention.

According to another aspect of the invention a method of maintaining or increasing the sucrose level in plant tissue comprises the step of transforming cells of the plant tissue with at least one gene construct of the invention.

According to another aspect of the invention a method of manipulating sucrose metabolism in a plant cell comprises the step of co-transforming the plant cell with each of the gene constructs of the invention.

The method may involve the alteration of sucrose metabolism in a plant or plant part containing stored sucrose.

The plant may be sugarcane.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be described in more detail, by way of example only, with reference to the accompanying drawings in which:

Figure 1 is the nucleotide sequence of the sugarcane PFP- β gene;

Figure 2 is a flow diagram of the steps involved in the isolation and characterization of the sugarcane PFP- β cDNA fragment;

Figure 3 is the nucleotide sequence of the 1209 base pair (bp) cDNA fragment containing the 3' end of the PFP- β gene; and

Figure 4 is a schematic representation of the genetic constructs, pUSPC 510 and pASPC 510, containing the 1209 bp PFP- β cDNA fragment in an untranslatable and an antisense form, respectively.

DETAILED DESCRIPTION OF THE INVENTION

A sugarcane leafroll cDNA library for PFP- β cDNA was screened as described below. Fragments of the PFP- β cDNA were isolated from the cDNA library. As an example, the constructs of a set of expression vectors containing a 1209 BP fragment of the 3'-end of the sugarcane PFP- β gene

will be described in detail. The fragment was used to construct vectors which were used to transform cells in a sugarcane callus. One of the vectors, pUSPC 510, contained the fragment in a sense orientation but lacked a translation initiation codon. The other vector, pASPC 510 contained the fragment in an antisense form. It was found that the isolated gene fragments could be used to regulate or manipulate the level of active PFP in the cells thereby manipulating sucrose metabolism in the cells.

Referring to Figure 2, an amplified PFP- β cDNA fragment was used as a probe to screen a sugarcane leafroll cDNA library for the PFP- β cDNA. Sequencing analysis was done to characterize the isolated clone. The sequence is shown in Figure 3. The insert of the clone was removed and used in the construction of two plant expression vectors. A promoter cassette with the CaMV 35S and the maize polyubiquitin (UBI) promoters was used in these constructs. In the first vector the fragment was cloned in the sense orientation, which is untranslatable because of the lack of a translation initiation codon. The fragment was cloned in the antisense orientation in the second vector. Schematic representations of the two expression vectors are shown in Figure 4.


Co-transformation with one of the expression vectors and a selectable marker gene of sugarcane callus was performed using a particle inflow gun and transformants were selected on geneticin-containing medium. Transformants were subsequently analyzed by measuring the PFP enzyme activity in the tissue with a standardised method (Botha *et al.* 1986). The PFP- β -fragment was excised from the library vector (pPFP 5) using the restriction enzymes Hinc III and Sma I. After purification, this fragment was cloned into the Sma I site of the expression vector DUBI 920.

References:

Botha, F. C., Small J.G.C., de Vries, C. (1986) Isolation and Characterisation of pyrophosphate: D-fructose-6-phosphate 1-phosphotransferase from cucumber seeds. Plant Cell Physiology 27: 1285-1295.

The content of this document is incorporated herein by reference.


Dated this 8th day of February 2000.



SPOOR AND FISHER

APPLICANT'S PATENT ATTORNEYS

10	20	30	40	50	60	70
ATGGCGGCGC	CGAGCGGACC	ATCACCTGGG	ACTGGGAGGT	TGGCGTCGGT	TTACAGCGAG	GTGCAGACGA
80	90	100	110	120	130	140
GCCGCTCCA	TCACGCGATC	CGGCTCCCCCT	CCGTCCTCTG	CTCCCAATTC	TCCCTCGTCG	ATGGACCTCC
150	160	170	180	190	200	210
CAGCTCAGCC	ACGGGGAACC	CGGATGAGAT	CGCGAAGCTG	TTCCCTAACT	TGTTTGGGCA	GCCGTCGGCG
220	230	240	250	260	270	280
ACATTGGTGC	CGGCCAAAGA	GGCGGTGGAG	GGGAAGGCGC	TGAAGGTCGG	GGTGGTGCTC	TCTGGTGGAC
290	300	310	320	330	340	350
AAGCACCCGG	TGGGCACAAT	GTGATCTGCG	GATCTTCTGA	TTTCTTGCG	AAACACGCAA	AGGGAAGCAC
360	370	380	390	400	410	420
AATGTATGGA	TTCAAAGGAG	GCCCAGCAGG	GGTGATGAAG	TGCAAGTACG	TCAAACCTCAA	TACCGATTTC
430	440	450	460	470	480	490
GTCTATCCCT	ACAGAAACCA	GGGTGGTTTT	GATATGATCT	GTAAGTGAAG	GGATAAGATT	GAAACACCAG
500	510	520	530	540	550	560
AGCAGTTTAA	GCAAGCCGAA	GATACAGCCA	ACAACTTGA	GTTGGACGGA	CTTGTTGTTA	TTGGACGGGA
570	580	590	600	610	620	630
CGATTCAAAT	ACTCATGCTT	GCCTCTTTGC	TGAATACTTC	AGGAGTAAAA	ATTTGAAAAC	CCGTGTCAAT
640	650	660	670	680	690	700
GGCAGCCCAA	AGACCATTGA	TGGTGATCTC	AAATGCAAAG	AGGTTCCAAC	CAGTTTTGGA	TTTGACACTG
710	720	730	740	750	760	770
CATGCAAGAT	CTATTCAGAA	ATGATTGSA	ATGTCATGAT	TGATGCCCGA	TCAACTGGAA	AATATTATCA
780	790	800	810	820	830	840
CTTTGTACGG	CTTATGGGGC	GTGCTGCTTC	TCACATTACA	TTGGGATGCG	CTTTGCAAAC	ACACCCCAAT
850	860	870	880	890	900	910
GCTGCACTCA	TTGGGGAAGA	GGTTGCTGCA	AAGAAGCAAA	CCCTTAAGAA	CGTCACAAAC	TACATTACTG
920	930	940	950	960	970	980
ATATCATCTG	CGAGCGTGCA	GATCTTGGTT	ACAACTATGG	TGTTATCCTT	ATACCAGAAG	GCCTGATTGA
990	1000	1010	1020	1030	1040	1050
TTTCATCCCA	GAGGTGCAGA	ATATCATTTG	TGAATTGAAT	GAAATTTTGG	CACATGATGT	TGTTGATGAG
1060	1070	1080	1090	1100	1110	1120
GCAGGGGCT	GGAAAAGCAA	GCTTCAGCCT	GAATCAAAGG	AGCTGTTTGA	GTTTTTGCCC	AAAACCTATTC
1130	1140	1150	1160	1170	1180	1190
AGGAGCAACT	TATGCTTGAA	AGGGGCCCCC	ATGGCAATGT	TCAGGTTGCA	AAAATTGAAA	CCGAGRAAAT
1200	1210	1220	1230	1240	1250	1260
GCTTATTAGC	ATGGTGGAAA	CTGAAGTGA	GAAGAGAAAA	GCAGAGGGGA	GATACTCTGC	ACATTTTCTA
1270	1280	1290	1300	1310	1320	1330
GGGCAAGCTC	ATTTCTTTGG	GTACGAAGGA	AGATGTGGCC	TTCTTACCAA	TTTTGATTCT	AACTATTGCT
1340	1350	1360	1370	1380	1390	1400
ATGCATTAGG	CTATGGGGCT	GCTGCCCTTC	TCCAAAGTGG	GAAGACAGGA	CTTATTTTCT	CGTTTGGCAA
1410	1420	1430	1440	1450	1460	1470
CCTTGCGGCT	CCAGTAGAAG	AATGGACTGT	TGGTGGAAAC	GCATTGACAT	CACTGATGGA	TGTGGAGAGG
1480	1490	1500	1510	1520	1530	1540
AGGCATGGCA	AGTTCAAGCC	AGTGATCGAG	AAGGCTATGG	TGGAACCTGA	TGCTGCACCT	TTCAAGAAAT
1550	1560	1570	1580	1590	1600	1610
ATGCATCAAT	GCGGGATGAG	TGGGCCACCA	AGAACAGATA	CATCAGCCCT	GGCCCCATCC	AGTTTCAAGT
1620	1630	1640	1650	1660	1670	1680
CCCTGGAAGT	GATGACTCGA	ACCACACTTT	GATGCTGGAA	CTCGGTGCTG	AGTTATAG...

Figure 1. The nucleotide sequence of the sugarcane PFP- β gene.





Figure 2. Flow diagram of the steps involved in the isolation and characterisation of sugarcane PFP- β cDNA and construction of expression vectors for the manipulation of sucrose metabolism in sugarcane.

10	20	30	40	50	60	70
GACGGTATCG	ATAAGCTTGA	TATCGAATTC	CGATTTAGCC	TCATACTGCT	TCTCACATT	CATTGGGATG
80	90	100	110	120	130	140
CGCTTTGCAA	ACACACCCCA	ATGCTGCACT	CATTGGGGAA	GAGGTTGCTG	CGAAGAAGCA	AACCCTTAAG
150	160	170	180	190	200	210
AACGTCACAA	ACTACATTAC	TGATATCATC	TGCAAGCGTG	CAGATCTTGG	TTACAACAT	GGGGTTATCC
220	230	240	250	260	270	280
TTATACCAGA	AGGCCTGATT	GATTTTCATC	CAGAGGTTCA	AAAACTCATC	GCAGAATTGA	ATGAAATTTT
290	300	310	320	330	340	350
GGCACATGAT	GTGGTTGATG	AGGCAGGGGC	CTGGAAAAGC	AAGCTTCAGC	CTGAATCAAA	GGAGCTGTTT
360	370	380	390	400	410	420
GAGTTTTTGC	CCAAAACAT	TCAGGAGCAA	CTTATGCTTG	AAAGGGGCCC	CCATGGCAAT	GTTCAGGTTG
430	440	450	460	470	480	490
CAAAAATTGA	AACCGAGAAA	ATGCTTATTA	GCATGGTGGA	AACTGAACTG	GAGAAGAGAA	AAGCAGAGGG
500	510	520	530	540	550	560
GAGATACTCT	GCACATTTCA	GAGGGCAAGC	TCATTTCTTT	GGGTACGAAG	GAAGATGTGG	CCTTCCTACC
570	580	590	600	610	620	630
AATTTTGATT	CTAACTATTG	CTATGCATTA	GGCTATGGTG	CTGGTGCCCT	TCTCCAAAGT	GGGAAGACAG
640	650	660	670	680	690	700
GACTTATTTT	ATCGGTTGGC	AACCTTGCGG	CTCCAGTAGA	AGAATGGACT	GTTGGTGGA	CAGCATTGAC
710	720	730	740	750	760	770
ATCACTGATG	GATGTTGAGA	GGAGGCATGG	CAAGTTCAAG	CCAGTGATCA	AGAAGGCTAT	GGTGGAACCT
780	790	800	810	820	830	840
GATGCTGCAC	CTTTCAAGAA	ATATGCATCA	ATGCGGGATG	AGTGGGCCAC	CAAGAACAGA	TACATCAGCC
850	860	870	880	890	900	910
CTGGCCCCAT	CCAGTTCAGT	GGCCCTGGAA	GTGATGACTC	GAACCACACT	TTGATGCTGG	AACCTCGGTG
920	930	940	950	960	970	980
TGAGTTATAG	AGATGCGTCC	TTTGCTTATT	TTTGTTTCTT	ACAGTTTGG	GAGTGGAGAC	TGGACACTGG
990	1000	1010	1020	1030	1040	1050
GTCTCCTGGA	GCAGCCTGCA	GTCTCCATAT	TGTGAATTGT	TTAATAAGAG	GTTTCGATGT	AGTTTTCTGC
1060	1070	1080	1090	1100	1110	1120
GTAGCGGACT	GGATGTAGCA	AATAAGAACT	GGTTTTAGCA	TTTTTTGTAT	GATTTACGCA	CCAACTGACT
1130	1140	1150	1160	1170	1180	1190
TGTCTTGTA	CCCTGATTCT	GTTCCACTGG	TTGCAATCTC	GTGAGAATGA	ACAAGTTGAT	ATGAGGCTAA
1200	1210	1220	1230	1240	1250	1260
ATCGGAATTC	CTGCAGCCC					

Figure 3. Nucleotide sequence of the 1209 bp fragment of the 3'-end of the sugarcane PFP- β -gene isolated from a cDNA library. The termination codon (TAG) is underlined (bp 918).






Figure 4. Schematic representation of the genetic constructs, a.) pUSPC 510 and b.) pASPC 510, containing 1209 bp of the PFP- β cDNA in the untranslatable (U) and antisense (A) forms respectively as examples of expression vectors containing sugarcane PFP- β sequences.